

COMPOSITION

Ibruxen Capsule: Each capsule contains Ibrutinib 140 mg INN.

INDICATIONS AND USAGE

Mantle Cell Lymphoma

Ibrutinib (**Ibruxen**) is indicated for the treatment of patients with Mantle Cell Lymphoma (MCL) who have received at least one prior therapy.

Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma

Ibrutinib (**Ibruxen**) is indicated for the treatment of adult patients with chronic lymphocytic leukemia (CLL)/ small lymphocytic lymphoma (SLL).

Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma with 17p deletion

Ibrutinib (**Ibruxen**) is indicated for the treatment of patients with chronic lymphocytic leukemia (CLL)/ small lymphocytic lymphoma (SLL) with 17p deletion.

Waldenstrom Macroglobulinemia (WM)

Ibrutinib (**Ibruxen**) is indicated for the treatment of patients with Waldenstrom Macroglobulinemia (WM).

Marginal Zone Lymphoma

Ibrutinib (**Ibruxen**) is indicated for the treatment of patients with marginal zone lymphoma (MZL) who require systemic therapy and have received at least one prior anti-CD20-based therapy.

Chronic Graft versus Host Disease

Ibrutinib (**Ibruxen**) is indicated for the treatment of adult patients with chronic graft-versus-host disease (cGVHD) after failure of one or more lines of systemic therapy

DOSAGE AND ADMINISTRATION

Dosing Guidelines

Administer Ibrutinib (**Ibruxen**) orally once daily at approximately the same time each day. Swallow the capsules whole with water. Do not open, break, or chew the capsules.

Recommended Dosage

Mantle Cell Lymphoma and Marginal Zone Lymphoma

The recommended dose of Ibrutinib (**Ibruxen**) for MCL and MZL is 560 mg (four 140 mg capsules) orally once daily until disease progression or unacceptable toxicity.

Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma and Waldenstrom Macroglobulinemia (WM)

The recommended dose of Ibrutinib (**Ibruxen**) for CLL/SLL and WM is 420 mg (three 140 mg capsules) orally once daily until disease progression or unacceptable toxicity. The recommended dose of Ibrutinib for CLL/SLL when used in combination with bendamustine and rituximab (administered every 28 days for up to 6 cycles) is 420 mg (three 140 mg capsules) orally once daily until disease progression or unacceptable toxicity.

Chronic Graft versus Host Disease

The recommended dose of Ibrutinib (**Ibruxen**) for cGVHD is 420 mg (three 140 mg capsules) orally once daily until cGVHD progression, recurrence of an underlying malignancy, or unacceptable toxicity. When a patient no longer requires therapy for the treatment of cGVHD, Ibrutinib should be discontinued considering the medical assessment of the individual patient.

Dose Modifications for Adverse Reactions

Interrupt Ibrutinib therapy for any Grade 3 or greater non-hematological toxicities, Grade 3 or greater neutropenia with infection or fever, or Grade 4 hematological toxicities. Once the symptoms of the toxicity have resolved to Grade 1 or baseline (recovery), Ibrutinib therapy may be reinitiated at the starting dose. If the toxicity reoccurs, reduce dose by one capsule (140 mg per day). A second reduction of dose by 140 mg may be considered as needed. If these toxicities persist or recur following two dose reductions, discontinue Ibrutinib.

Toxicity Occurrence	MCL and MZL Dose Modification After Recovery Starting Dose=560mg	CLL/SLL and WM Dose Modification After Recovery Starting Dose=420mg
First	Restart at 560 mg daily	Restart at 420 mg daily
Second	Restart at 420 mg daily	Restart at 280 mg daily
Third	Restart at 280 mg daily	Restart at 140 mg daily
Fourth	Discontinue Ibrutinib	Discontinue Ibrutinib

Dose Modifications for Use with CYP3A Inhibitors

Avoid co-administration with strong or moderate CYP3A inhibitors and consider alternative agents with less CYP3A inhibition. Concomitant use of strong CYP3A inhibitors which would be taken chronically (e.g., ritonavir, indinavir, nelfinavir, saquinavir, boceprevir, telaprevir, nefazodone) is not recommended. For short-term use (treatment for 7 days or less) of strong CYP3A inhibitors (e.g., antifungals and antibiotics) consider interrupting Ibrutinib therapy until the CYP3A inhibitor is no longer needed. Reduce Ibrutinib dose to 140 mg if a moderate CYP3A inhibitor must be used (e.g., fluconazole, darunavir, erythromycin, diltiazem, atazanavir, aprepitant, amprenavir, fosamprenavir, crizotinib, imatinib, verapamil, and ciprofloxacin). Patients taking concomitant strong or moderate CYP3A inhibitors should be monitored more closely for signs of Ibrutinib toxicity.

Dose Modifications for Use in Hepatic Impairment

For patients with mild liver impairment (Child-Pugh class A), the recommended dose is 140 mg daily (one capsule). Avoid the use of Ibrutinib in patients with moderate or severe hepatic impairment (Child-Pugh classes Band C).

Missed Dose

If a dose of Ibrutinib is not taken at the scheduled time, it can be taken as soon as possible on the same day with a return to the normal schedule the following day. Extra capsules of Ibrutinib should not be taken to make up for the missed dose.

CONTRAINDICATIONS

Hypersensitivity to Ibrutinib or to any of the excipients.

WARNINGS AND PRECAUTIONS

Hemorrhage

Fatal bleeding events have occurred in patients treated with

Ibrutinib. Grade 3 or higher bleeding events (intracranial hemorrhage [including subdural hematoma], gastrointestinal bleeding, hematuria, and post procedural hemorrhage) have occurred in up to 6% of patients. Bleeding events of any grade, including bruising and petechiae, occurred in approximately half of patients treated with Ibrutinib. The mechanism for the bleeding events is not well understood. Ibrutinib may increase the risk of hemorrhage in patients receiving antiplatelet or anticoagulant therapies and patients should be monitored for signs of bleeding. Consider the benefit-risk of withholding Ibrutinib for at least 3 to 7 days pre and post-surgery depending upon the type of surgery and the risk of bleeding.

Infections

Fatal and non-fatal infections (including bacterial, viral, or fungal) have occurred with Ibrutinib therapy. Grade 3 or greater infections occurred in 14% to 29% of patients. Cases of progressive multifocal leukoencephalopathy (PML) and Pneumocystis jirovecii pneumonia (PJP) have occurred in patients treated with Ibrutinib. Consider prophylaxis according to standard of care in patients who are at increased risk for opportunistic infections. Monitor and evaluate patients for fever and infections and treat appropriately.

Cytopenias

Treatment-emergent Grade 3 or 4 cytopenias including neutropenia (range, 13 to 29%), thrombocytopenia (range, 5 to 17%), and anemia (range, 0 to 13%) based on laboratory measurements occurred in patients with B-cell malignancies treated with single agent Ibrutinib. Monitor complete blood counts monthly.

Cardiac Arrhythmias

Fatal and serious cardiac arrhythmias have occurred with Ibrutinib therapy. Grade 3 or greater ventricular tachyarrhythmias occurred in 0 to 1% of patients, and Grade 3 or greater atrial fibrillation and atrial flutter occurred in 0 to 6% of patients. These events have occurred particularly in patients with cardiac risk factors, hypertension, acute infections, and a previous history of cardiac arrhythmias. Periodically monitor patients clinically for cardiac arrhythmias. Obtain an ECG for patients who develop arrhythmic symptoms (e.g., palpitations, lightheadedness, syncope, chest pain) or new onset dyspnea. Manage cardiac arrhythmias appropriately, and if it persists, consider the risks and benefits of Ibrutinib treatment and follow dose modification guidelines.

Hypertension

Hypertension (range, 6 to 17%) has occurred in patients treated with Ibrutinib with a median time to onset of 4.6 months (range, 0.03 to 22 months). Monitor patients for new onset hypertension or hypertension that is not adequately controlled after starting Ibrutinib. Adjust existing anti-hypertensive medications and/or initiate anti-hypertensive treatment as appropriate.

Second Primary Malignancies

Other malignancies (range, 3 to 16%) including non-skin carcinomas (range, 1 to 4%) have occurred in patients treated with Ibrutinib. The most frequent second primary malignancy was non-melanoma skin cancer (range, 2 to 13%).

Tumor Lysis Syndrome

Tumor lysis syndrome has been infrequently reported with Ibrutinib therapy. Assess the baseline risk (e.g., high tumor burden) and take appropriate precautions. Monitor patients closely and treat as appropriate.

Embryo-Fetal Toxicity

Based on findings in animals, Ibrutinib can cause fetal harm when administered to a pregnant woman. Administration of Ibrutinib to pregnant rats and rabbits during the period of organogenesis caused embryo-fetal toxicity including malformations at exposures that were 2-20 times higher than those reported in patients with hematologic malignancies. Advise women to avoid becoming pregnant while taking Ibrutinib and for 1 month after cessation of therapy. If this drug is used during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to a fetus.

SIDE EFFECTS

The following adverse reactions are discussed in more detail in other sections of the labeling: Hemorrhage, Infections, Cytopenias, Atrial Fibrillation, Hypertension, Second Primary Malignancies and Tumor Lysis Syndrome.

The most commonly occurring adverse reactions (≥ 20%) were thrombocytopenia, diarrhea, neutropenia, anemia, fatigue, musculoskeletal pain, peripheral edema, upper respiratory tract infection, nausea, bruising, dyspnea, constipation, rash, abdominal pain, vomiting and decreased appetite. The most common Grade 3 or 4 non-hematological adverse reactions (≥ 5%) were pneumonia, abdominal pain, atrial fibrillation, diarrhea, fatigue, and skin infections.

DRUG INTERACTIONS

Effect of CYP3A Inhibitors on Ibrutinib

Ibrutinib is primarily metabolized by cytochrome P450 enzyme 3A (CYP3A). In healthy volunteers, co-administration of ketoconazole, a strong CYP3A inhibitor, increased Cmax and AUC of Ibrutinib by 29- and 24-fold, respectively. The highest Ibrutinib dose evaluated in clinical trials was 12.5 mg/kg (actual doses of 840–1400 mg) given for 28 days with single dose AUC values of 1445 ± 869 ng·hr/mL which is approximately 50% greater than steady state exposures seen at the highest indicated dose (560 mg).

Avoid concomitant administration of Ibrutinib with strong or moderate inhibitors of CYP3A. For strong CYP3A inhibitors used short-term (e.g., antifungals and antibiotics for 7 days or less, e.g., ketoconazole, itraconazole, voriconazole, posaconazole, clarithromycin, telithromycin) consider interrupting Ibrutinib therapy during the duration of inhibitor use. Avoid strong CYP3A inhibitors that are needed chronically. If a moderate CYP3A inhibitor must be used, reduce the Ibrutinib dose. Patients taking concomitant strong or moderate CYP3A4 inhibitors should be monitored more closely for signs of Ibrutinib toxicity. Avoid grape fruit and Seville oranges during Ibrutinib treatment, as these contain moderate inhibitors of CYP3A.

Effect of CYP3A Inducers on Ibrutinib

Administration of Ibrutinib with rifampin, a strong CYP3A inducer, decreased Ibrutinib Cmax and AUC by approximately 13 & 10 fold, respectively. Avoid concomitant use of strong CYP3A inducers (e.g., carbamazepine, rifampin, phenytoin, and St. John'sWort). Consider alternative agents with less CYP3A induction.